

HIGHER ANALOGUES OF THE DISCRETE-TIME TODA EQUATION AND THE QUOTIENT-DIFFERENCE ALGORITHM

PAUL E. SPICER, FRANK W. NIJHOFF, AND PETER H. VAN DER KAMP

ABSTRACT. The discrete-time Toda equation arises as a universal equation for the relevant Hankel determinants associated with one-variable orthogonal polynomials through the mechanism of adjacency, which amounts to the inclusion of shifted weight functions in the orthogonality condition. In this paper we extend this mechanism to a new class of two-variable orthogonal polynomials where the variables are related via an elliptic curve. This leads to a ‘Higher order Analogue of the Discrete-time Toda’ (HADT) equation for the associated Hankel determinants, together with its Lax pair, which is derived from the relevant recurrence relations for the orthogonal polynomials. In a similar way as the quotient-difference (QD) algorithm is related to the discrete-time Toda equation, a novel quotient-quotient-difference (QQD) scheme is presented for the HADT equation. We show that for both the HADT equation and the QQD scheme, there exists well-posed s -periodic initial value problems, for almost all $s \in \mathbb{Z}^2$. From the Lax-pairs we furthermore derive invariants for corresponding reductions to dynamical mappings for some explicit examples.

1. INTRODUCTION

The discrete-time Toda equation, i.e. the time-discretized version of the usual Toda chain, which is given by the following partial difference equation (P Δ E)

$$\tau_{l-1,m-1}\tau_{l+1,m+1} = \tau_{l+1,m-1}\tau_{l-1,m+1} - \tau_{l,m}^2, \quad (1.1)$$

plays an important role in many areas of mathematical physics. Originally derived, albeit in a slightly different form, by Hirota, [23], it is the equation that generalizes to the space-time lattice (here indicated by the discrete independent variables $(l, m) \in \mathbb{Z}^2$) the Toda chain in such a way that the prominent integrability features of the latter equation (such as the existence of soliton solutions, existence of an underlying linear problem – its Lax pair– etc.) are retained. This P Δ E has also appeared in physics, namely as the nonlinear equation governing the spin-spin correlation functions of the two-dimensional Ising model, cf. [34, 40]. The connection we investigate and generalize in this paper, is the one with the theory of formal orthogonal polynomials and Padé approximants, where the equation (1.1) is connected to the well known quotient-difference (QD) algorithm

$$\begin{aligned} e_{l,m+1} + q_{l+1,m+1} &= q_{l+1,m} + e_{l+1,m}, \\ e_{l,m+1}q_{l,m+1} &= q_{l+1,m}e_{l,m}. \end{aligned} \quad (1.2)$$

The theory of formal orthogonal polynomials is a subject central to modern numerical analysis, where orthogonalities are investigated not so much on the particular problems associated with specific weight functions (such as the moment problem, or the problem of the behaviour of the zeroes of the polynomial) but rather how general properties arising from orthogonality can be implemented in connection with numerical algorithms. In this area of research general constructions such as

those of vector Padé approximants, adjacent orthogonal families (where connected sequences of orthogonality functionals are postulated), and associated convergence acceleration algorithms, and factorisation methods are investigated, cf. e.g. [8, 15]. In this context, the QD algorithm emerges as a prominent method to locate the zeroes of analytic functions or to compute convergence factors for asymptotic expansions through formal power series, using a finite-difference scheme and continued fractions, cf. [20]. It was pointed out in [39] that both the QD algorithm and certain convergence algorithms, [18], are intimately related to integrable discrete systems. In fact, the famous ϵ -algorithm of Wynn, [47], surprisingly turns out to be identical to the lattice potential Korteweg-de Vries (KdV) equation, a well-known exactly integrable P Δ E, thus allowing us to interpret the numerical algorithm as a symplectic dynamical system with rich behaviour from the point of view of analytical solutions. Similarly, the famous “missing identity of Frobenius”, [17], in the theory of Padé approximants can be regarded as an exactly solvable lattice system closely related to discretisations of the KdV equation and intimately connected to the Toda lattice.

In this paper we introduce a novel integrable P Δ E:

$$\begin{aligned} & \sigma_{l+1,m-2}\sigma_{l-1,m+1} (\sigma_{l,m+2}\sigma_{l,m-1} - \sigma_{l,m}\sigma_{l,m+1}) \\ = & \sigma_{l,m-1}\sigma_{l-1,m+2} (\sigma_{l-1,m+1}\sigma_{l+2,m-2} - \sigma_{l,m}\sigma_{l+1,m-1}) \\ + & \sigma_{l,m+1}\sigma_{l+1,m-1} (\sigma_{l,m-2}\sigma_{l-1,m+2} - \sigma_{l+1,m-2}\sigma_{l-2,m+2}), \end{aligned} \quad (1.3)$$

which we argue can be regarded as a Higher order Analogue of the Discrete-time Toda equation and therefore name it the HADT equation. It is defined on a stencil of 11 points in the lattice as depicted in Figure 1.




FIGURE 1. The stencil of the HADT equation (1.3)

The derivation of the HADT equation is parallel to the way in which the discrete-time Toda equation emerges in the theory of orthogonal polynomials: in fact, we introduce a family of two-variable orthogonal polynomials restricted by an elliptic curve and exploit the recurrence structure for the relevant Hankel determinants. The general problem of orthogonal polynomials on algebraic curves was discussed in the monograph by Suetin, [44, Chapter 7], and has also been addressed in the context of the study of formal orthogonalities, cf. e.g. [11]. The construction of the associated two-variable orthogonal polynomials on elliptic curve is pursued in [43], whereas in this paper we will concentrate on the HADT equation (1.3) itself and its reductions. Nevertheless, we will present the derivation from the adjacency structure of elliptic two-variable orthogonal polynomials, as it produces not only the nonlinear equation itself but also its Lax pair. Furthermore, the

recurrence structure of the relevant Hankel determinants also yields a novel type of QD formalism, namely the following so-called quotient-quotient-difference (QQD) scheme:

$$\begin{aligned} u_{l+2,m} + v_{l+1,m} + w_{l+1,m+1} &= u_{l,m+3} + v_{l+1,m+1} + w_{l+1,m}, \\ u_{l,m+3}v_{l,m+1} &= v_{l+1,m}u_{l+1,m}, \\ u_{l,m+3}w_{l,m+1} &= w_{l+1,m+1}u_{l+1,m+1}, \end{aligned} \tag{1.4}$$

which we believe may have future applications in numerical analysis.

Once we have obtained the HADT equation, the QQD scheme and their Lax pairs, we address the issue of periodic reductions. A periodic solution of a (system of) lattice equation(s) is a solution which satisfies $\mathbf{u}_{l,m} = \mathbf{u}_{l+s_1,m+s_2}$ for some $s_1, s_2 \in \mathbb{Z}$. By imposing periodicity a lattice equation reduces to a (system of) ordinary difference equations, or, equivalently, a mapping. The so-called *staircase method* utilizes the Lax-pair of the lattice equation to construct integrals for the mapping. The method was introduced in [38], where it was applied to the KdV equation and to a mixed modified KdV-Toda equation, taking $s_1 = s_2$ and $s_1 = s_2 + 1$. In [41] a two-parameter family of periodic reductions was studied, with s_1 and s_2 being co-prime integers. Recently, in [25, 42] a unified picture for periodic reductions, with nonzero $\mathbf{s} = (s_1, s_2) \in \mathbb{Z}^2$ has been provided. In [42] it was shown how, under such a periodicity condition, any lattice equation $f(u_{l,m}, \dots) = 0$ reduces to a system of q ordinary difference equations $f(v_n^p, \dots) = 0$, $p = 0, \dots, r-1$, where r is the greatest common divisor of s_1 and s_2 . Also it was proved that the monodromy matrix, denoted \mathcal{L} , is one of the Lax matrices for the reduction. In [25] it was shown, for any given scalar lattice equation on some arbitrary stencil of lattice points, there exists a well-posed, or *nearly well-posed*, \mathbf{s} -periodic initial value problem, for all nonzero $\mathbf{s} \in \mathbb{Z}^2$. Thus, the trace of the monodromy matrix provides integrals for any periodic reduction of any integrable scalar lattice equation, and it is expected that the same is true for systems of lattice equations, see [25, 27]. In this paper we generalize the approach of [25] and develop a systematic method to construct well-posed periodic initial value problems for systems of lattice equations. This method is applied to the QQD scheme as well as to an intermediate system of PΔEs, cf. equation (3.19).

On general principles, one expects periodic reductions of integrable lattice equations to lead to integrable maps in a precise sense, namely to complete integrability in the sense of Liouville-Arnold [12, 46]. This requires, in addition to the existence of a sufficient number of integrals, some global properties as well the existence of a symplectic structure which is preserved by the discrete dynamics, and with respect to which the integrals are in involution. The latter are issues which we are not addressing in the current paper, where we have developed the basic structures and classified the consistent reductions from the lattice equation together with the explicit form of the integrals. In order to pursue the next step, we first need to establish appropriate canonical variables which are not necessarily the variables directly given on the vertices of the lattice. In fact, typically some *reduced variables*, i.e. suitable combinations of the variables on the vertices, need to be introduced, leading to a further dimensional reduction of the mappings. Furthermore, there is the problem of counting the number of independent integrals to ensure that in principle a 1-1 map to the relevant number of degrees of freedoms exist for the mappings concerned.

For the reductions we obtained from the HADT equation, as well as those from the QQD scheme, we are able to find a sufficient number of integrals, that is, equal to half the dimensionality (i.e. number of components) of the mapping. For the reductions obtained from the intermediate system (3.19) this, however, is not the case. In [27] a systematic method was presented, which exploits symmetries of the (system of) lattice equation(s) to reduce the dimension of the mappings. For all cases that were considered in [27], the staircase method provides a sufficient number of integrals for the dimensionally reduced mappings to be in principle completely integrable.

In this paper we encounter one case, the $\mathbf{s} = (2, -3)$ reduction of system (3.19), where this approach yields a five-dimensional mapping with only two integrals. We have not been able yet to find an extra integral, nor an extra symmetry, but we are confident that the corresponding mapping is indeed integrable, as we have verified that it has quadratic growth. We note that polynomial growth of the degrees of iterates of the mapping means vanishing algebraic entropy (a concept introduced by Viallet and collaborators [6, 16]) and this corresponds to low complexity (which characterises integrable maps [1, 46]). Examples show that degree growth is a better indication of integrability than singularity confinement [21, 22], cf. the discussion in [37]. We refer to [26] for an elementary method to actually prove polynomial growth.

2. THE DISCRETE-TIME TODA EQUATION AND THE QD ALGORITHM FROM ORTHOGONAL POLYNOMIALS

We present here the derivation of the bilinear discrete-time Toda equation using formal orthogonal polynomials. Recurrence relations for adjacent one-variable orthogonal polynomials provide a Lax-pair for this equation, as well as for the associated QD algorithm. The results presented here are known, see for example [39], however they outline the main idea which we employ in the setting of two-variable elliptic orthogonal polynomials in the next section.

Let P_n be a family of formal orthogonal polynomials defined with respect to a weight function $w(x)$ on a curve Γ in the complex plane or, in other words, with respect to the linear functional c_n such that

$$\mathcal{L}(e_n) = c_n = \int_{\Gamma} x^n w(x) dx \quad i = 0, 1, \dots \quad (2.1)$$

where e_n are the monomials. The P_n 's, which are assumed to be monic, satisfy a three term recurrence relation of the form

$$xP_n = P_{n+1} + S_n P_n + R_n P_{n-1} \quad (2.2)$$

(where S_n and R_n are finite) and in order to insure their existence we assume that the corresponding Hankel determinants are non-zero.

We introduce a family of adjacent orthogonal polynomials (which are orthogonal with respect to a shifted weight $x^m w(x)$) [5]:

$$P_n^{(m)}(x) \equiv \frac{1}{\Delta_{n-1}^{(m)}} \begin{vmatrix} c_m & \dots & \dots & c_{n+m} \\ \vdots & & & \vdots \\ c_{n+m-1} & & & c_{2n+m-1} \\ 1 & \dots & \dots & x^n \end{vmatrix}, \quad (2.3a)$$

We will refer to equation (2.8) as **the discrete-time Toda equation**. Indeed, substituting $\Delta_n^{(m)} = \tau_{-2n-m-1, m-1}$ and taking $l = -2n - m$ as new independent variable yields the more standard form of this equation (1.1). Thus the shadows of integrability already appear in the underlying structure of the standard theory of orthogonal polynomials.

2.1. A Lax pair for the QD algorithm and the discrete-time Toda equation. We now change perspective and view equation (2.8) as an integrable lattice equation, as opposed to an identity for Hankel determinants (2.3b). To this end we write $\Delta_n^{(m)} = \tau_{n,m}$, and similarly $V_n^{(m)} = v_{n,m}$, $W_n^{(m)} = w_{n,m}$. The variable x will play the role of spectral parameter and will be denoted $x = \lambda$.

The relations (2.5) constitute a Lax-pair for equation (2.8). Working with the fields v, w we first derive a related quotient-difference system. Let $\Psi_{n,m} = (P_n^{(m)}, P_n^{(m+1)})$, then using the recurrence relations we derive

$$\Psi_{n+1,m} = L_{n,m} \Psi_{n,m} \quad \text{and} \quad \Psi_{n,m+1} = M_{n,m} \Psi_{n,m},$$

with

$$L_{n,m} = \begin{pmatrix} -v_{n,m} & \lambda \\ -v_{n,m} & \lambda + w_{n,m} - v_{n,m+1} \end{pmatrix}, \quad (2.9a)$$

$$M_{n,m} = \begin{pmatrix} 0 & 1 \\ -\lambda^{-1}v_{n,m} & 1 + \lambda^{-1}w_{n,m} \end{pmatrix}. \quad (2.9b)$$

The compatibility of the two linear systems is then equivalent to the discrete Lax equation:

$$\begin{aligned} 0 &= L_{n,m+1}M_{n,m} - M_{n+1,m}L_{n,m} \\ &= \begin{pmatrix} 0 & 0 \\ -\lambda^{-1}v_{n,m}E & E + \lambda^{-1}(w_{n,m}(v_{n+1,m} + E) - v_{n,m+1}w_{n+1,m}) \end{pmatrix}, \end{aligned}$$

where

$$E = v_{n,m+2} - v_{n+1,m} + w_{n+1,m} - w_{n,m+1}.$$

Thus L, M provide a Lax pair for the quotient-difference scheme

$$v_{n,m+2} + w_{n+1,m} = v_{n+1,m} + w_{n,m+1}, \quad (2.10a)$$

$$w_{n,m}v_{n+1,m} = v_{n,m+1}w_{n+1,m}. \quad (2.10b)$$

This scheme is related to the (more standard) QD algorithm (1.2), by $q_{n,m} = v_{n,m}$, $e_{n,m} = v_{n,m+1} - w_{n,m}$.

Upon substitution of (2.6) into (2.10a) this equation is a consequence of (2.8), whereas substitution of (2.6) into (2.10b) turns into an identity. To find a *good* Lax-pair for equation (2.8), one substitutes equations (2.6) into the above Lax matrices, and one uses (2.8) to simplify

$$e_{n,m} = v_{n,m+1} - w_{n,m} = \frac{\tau_{n-1,m+1}\tau_{n+1,m}}{\tau_{n-1,m+1}\tau_{n,m}}.$$

Using the same simplification, equation (2.10a) can be written

$$\begin{aligned} 0 &= e_{n,m+1} + w_{n+1,m} - v_{n+1,m} \\ &= \frac{\tau_{n+1,m+1}}{\tau_{n,m+2}\tau_{n,m+1}\tau_{n+1,m}} \left(\tau_{n+1,m}\tau_{n-1,m+2} - \tau_{n,m+2}\tau_{n,m} + (\tau_{n,m+1})^2 \right), \end{aligned}$$

where one recognizes the discrete-time Toda equation (2.8).

3. THE HADT EQUATION AND A QQD SCHEME FROM ELLIPTIC ORTHOGONAL POLYNOMIALS

In this section we derive the HADT equation. We follow a similar route as in the previous section, except that here we consider two-variable orthogonal polynomials, where the variables are restricted by the condition that they form the coordinates of an elliptic curve. The recurrence relations for these elliptic orthogonal polynomials yield both a Lax pair and the QQD scheme.

3.1. Two-variable elliptic orthogonal polynomials. As a starting point for our construction we introduce the sequence of elementary monomials [43], associated with a class of two variable orthogonal polynomials. We consider a sequence of monomials where the x and y are given different weight, namely $x \sim 2$ and $y \sim 3$:

$$e_0 = 1, \quad e_2 = x, \quad e_3 = y, \quad e_4 = x^2, \quad e_5 = xy, \quad e_6 = x^3, \quad \dots$$

or, in general:

$$e_0(x, y) = 1, \quad e_{2k}(x, y) = x^k, \quad e_{2k+1}(x, y) = x^{k-1}y, \quad k = 1, 2, \dots$$

In comparison with the polynomials of Krall and Scheffer [29], their two-variable orthogonal polynomials consist of $x \sim 1$ and $y \sim 1$ of equal weight. The reason for choosing different weights is the variables x and y are related through a Weierstrass elliptic curve. There is a famous connection between the monomial sequences of the form given above and the genus of the corresponding curve, given by the *Weierstrass gap sequence* [3], which states that the latter equals the number of omissions in the sequence. Since, in the present case we have a curve of genus 1, this accounts for the omission of the monomial e_1 . We use the sequence as our basis for the expansion of a new class of two-variable orthogonal polynomials taking the form:

$$P_k(x, y) = \sum_{j=0}^k p_j^{(k)} e_j(x, y), \quad (3.1)$$

which are *monic* if the leading coefficient $p_k^{(k)} = 1$.

In the spirit of the formal approach to orthogonal polynomials, cf. e.g. [10, 8], we assume that a bilinear form \langle, \rangle exists and can be derived from a linear functional \mathcal{L} and consequently we can define the associated *moments* by

$$c_k = \mathcal{L}(e_k). \quad (3.2)$$

The assumption of the existence of an inner product \langle, \rangle on the space \mathcal{V} spanned by the monomials e_k , is such that

$$\langle xP, Q \rangle = \langle P, xQ \rangle, \quad ,$$

for any two elements $P, Q \in \mathcal{V}$.

Under the assumption of orthogonality and using the standard Gram-Schmidt orthogonalisation (through the use of Cramer's rule), leads to the following expression for the adjacent elliptic orthogonal polynomials:

$$P_k^{(l)}(x, y) \equiv \begin{vmatrix} \langle \mathbf{e}_l, \mathbf{e}_0 \rangle & \langle \mathbf{e}_l, \mathbf{e}_2 \rangle & \cdots & \cdots & \langle \mathbf{e}_l, \mathbf{e}_k \rangle \\ \langle \mathbf{e}_{l+1}, \mathbf{e}_0 \rangle & \langle \mathbf{e}_{l+1}, \mathbf{e}_2 \rangle & \cdots & \cdots & \langle \mathbf{e}_{l+1}, \mathbf{e}_k \rangle \\ \vdots & \vdots & & & \vdots \\ \vdots & \vdots & & & \vdots \\ \langle \mathbf{e}_{l+k-2}, \mathbf{e}_0 \rangle & \langle \mathbf{e}_{l+k-2}, \mathbf{e}_2 \rangle & \cdots & \cdots & \langle \mathbf{e}_{l+k-2}, \mathbf{e}_k \rangle \\ \mathbf{e}_0 & \mathbf{e}_2 & \cdots & \cdots & \mathbf{e}_k \end{vmatrix} / \Delta_{k-1}^{(l)}, \quad l \neq 0, \quad (3.3)$$

together with the corresponding Hankel determinant:

$$\Delta_k^{(l)} = \begin{vmatrix} \langle \mathbf{e}_l, \mathbf{e}_0 \rangle & \langle \mathbf{e}_l, \mathbf{e}_2 \rangle & \cdots & \cdots & \langle \mathbf{e}_l, \mathbf{e}_k \rangle \\ \langle \mathbf{e}_{l+1}, \mathbf{e}_0 \rangle & \langle \mathbf{e}_{l+1}, \mathbf{e}_2 \rangle & \cdots & \cdots & \langle \mathbf{e}_{l+1}, \mathbf{e}_k \rangle \\ \vdots & \vdots & & & \vdots \\ \vdots & \vdots & & & \vdots \\ \langle \mathbf{e}_{l+k-1}, \mathbf{e}_0 \rangle & \langle \mathbf{e}_{l+k-1}, \mathbf{e}_2 \rangle & \cdots & \cdots & \langle \mathbf{e}_{l+k-1}, \mathbf{e}_k \rangle \end{vmatrix}, \quad l \neq 0, \quad (3.4)$$

where for $l = 0$:

$$P_k^{(0)}(x, y) \equiv \begin{vmatrix} \langle \mathbf{e}_0, \mathbf{e}_0 \rangle & \langle \mathbf{e}_0, \mathbf{e}_2 \rangle & \cdots & \cdots & \langle \mathbf{e}_0, \mathbf{e}_k \rangle \\ \langle \mathbf{e}_2, \mathbf{e}_0 \rangle & \langle \mathbf{e}_2, \mathbf{e}_2 \rangle & \cdots & \cdots & \langle \mathbf{e}_2, \mathbf{e}_k \rangle \\ \vdots & \vdots & & & \vdots \\ \vdots & \vdots & & & \vdots \\ \langle \mathbf{e}_{k-1}, \mathbf{e}_0 \rangle & \langle \mathbf{e}_{k-1}, \mathbf{e}_2 \rangle & \cdots & \cdots & \langle \mathbf{e}_{k-1}, \mathbf{e}_k \rangle \\ \mathbf{e}_0 & \mathbf{e}_2 & \cdots & \cdots & \mathbf{e}_k \end{vmatrix} / \Delta_{k-1}^{(0)}, \quad (3.5)$$

with

$$\Delta_k^{(0)} = \begin{vmatrix} \langle \mathbf{e}_0, \mathbf{e}_0 \rangle & \langle \mathbf{e}_0, \mathbf{e}_2 \rangle & \cdots & \cdots & \langle \mathbf{e}_0, \mathbf{e}_k \rangle \\ \langle \mathbf{e}_2, \mathbf{e}_0 \rangle & \langle \mathbf{e}_2, \mathbf{e}_2 \rangle & \cdots & \cdots & \langle \mathbf{e}_2, \mathbf{e}_k \rangle \\ \vdots & \vdots & & & \vdots \\ \vdots & \vdots & & & \vdots \\ \langle \mathbf{e}_k, \mathbf{e}_0 \rangle & \langle \mathbf{e}_k, \mathbf{e}_2 \rangle & \cdots & \cdots & \langle \mathbf{e}_k, \mathbf{e}_k \rangle \end{vmatrix}, \quad (3.6)$$

In addition to the polynomials (3.3), we also need to introduce the polynomials:

$$Q_k^{(l)}(x, y) \equiv \begin{vmatrix} \langle \mathbf{e}_l, \mathbf{e}_0 \rangle & \langle \mathbf{e}_l, \mathbf{e}_2 \rangle & \cdots & \cdots & \langle \mathbf{e}_l, \mathbf{e}_k \rangle \\ \langle \mathbf{e}_{l+2}, \mathbf{e}_0 \rangle & \langle \mathbf{e}_{l+2}, \mathbf{e}_2 \rangle & \cdots & \cdots & \langle \mathbf{e}_{l+2}, \mathbf{e}_k \rangle \\ \vdots & \vdots & & & \vdots \\ \vdots & \vdots & & & \vdots \\ \langle \mathbf{e}_{l+k-1}, \mathbf{e}_0 \rangle & \langle \mathbf{e}_{l+k-1}, \mathbf{e}_2 \rangle & \cdots & \cdots & \langle \mathbf{e}_{l+k-1}, \mathbf{e}_k \rangle \\ \mathbf{e}_0 & \mathbf{e}_2 & \cdots & \cdots & \mathbf{e}_k \end{vmatrix} / \Theta_{k-1}^{(l)}, \quad (3.7)$$

together with its corresponding Hankel determinant:

$$\Theta_k^{(l)} = \begin{vmatrix} \langle \mathbf{e}_l, \mathbf{e}_0 \rangle & \langle \mathbf{e}_l, \mathbf{e}_2 \rangle & \cdots & \cdots & \langle \mathbf{e}_l, \mathbf{e}_k \rangle \\ \langle \mathbf{e}_{l+2}, \mathbf{e}_0 \rangle & \langle \mathbf{e}_{l+2}, \mathbf{e}_2 \rangle & \cdots & \cdots & \langle \mathbf{e}_{l+2}, \mathbf{e}_k \rangle \\ \vdots & \vdots & & & \vdots \\ \vdots & \vdots & & & \vdots \\ \langle \mathbf{e}_{l+k}, \mathbf{e}_0 \rangle & \langle \mathbf{e}_{l+k}, \mathbf{e}_2 \rangle & \cdots & \cdots & \langle \mathbf{e}_{l+k}, \mathbf{e}_k \rangle \end{vmatrix}, \quad (3.8)$$

noting that

$$Q_k^{(0)} = P_k^{(0)}, \quad \Theta_k^{(0)} = \Delta_k^{(0)}.$$

Remark: We note that for $l \neq 0, 1$ the polynomials $P_k^{(l)}$ do *not* form an orthogonal family. In fact, from the determinantal definition (3.3) we immediately observe that

$$\langle \mathbf{e}_l, P_k^{(l)} \rangle = \langle \mathbf{e}_{l+1}, P_k^{(l)} \rangle = \cdots = \langle \mathbf{e}_{l+k-2}, P_k^{(l)} \rangle = 0, \quad \langle \mathbf{e}_{l+k-1}, P_k^{(l)} \rangle = \frac{\Delta_k^{(l)}}{\Delta_{k-1}^{(l)}},$$

whereas

$$\langle \mathbf{e}_{l-1}, P_k^{(l)} \rangle = (-1)^{k-1} \frac{\Delta_k^{(l-1)}}{\Delta_{k-1}^{(l)}}.$$

We now proceed using determinantal identities of Sylvester type (Appendix A) to derive relations between the polynomials $P_k^{(l)}$ and the Hankel determinants.

3.2. Recurrence relations in $P_k^{(l)}$ and $Q_k^{(l)}$. Using a 3 row/column Sylvester identity we find it is possible to find x -recurrence relations and linear relations in $P_k^{(l)}$ and $Q_k^{(l)}$. To achieve the former it is necessary to fix the columns, so that $\mathbf{e}_0 = 1$ and $\mathbf{e}_3 = y$ are removed from the determinant and the position of the row removal is dependent on restricting the introduction of new objects. Hence we apply the following cutting of three rows and columns (A.6) to the determinant for $P_k^{(l)}$:

(where the small red line indicates a space between the first column and the third).

We are led to the recurrence relation

$$P_k^{(l)} = xP_{k-2}^{(l+3)} - V_{k-2}^{(l)}P_{k-1}^{(l)} + W_{k-2}^{(l)}P_{k-1}^{(l+1)}, \quad l \neq 0, 1, \quad (3.9)$$

where

$$V_k^{(l)} = \frac{\Delta_k^{(l)} \Delta_k^{(l+3)}}{\Delta_{k+1}^{(l)} \Delta_{k-1}^{(l+3)}}, \quad W_k^{(l)} = \frac{\Delta_k^{(l+1)} \Delta_k^{(l+2)}}{\Delta_{k+1}^{(l)} \Delta_{k-1}^{(l+3)}}. \quad (3.10)$$

which can be expressed in a simpler way using

$$\begin{aligned} X_k^{(l)} &= \Gamma_k^{(l)} - \Gamma_{k-1}^{(l)}, \\ Y_k^{(l)} &= \Gamma_{k-2}^{(l+2)} - \Gamma_{k-1}^{(l)}, \end{aligned}$$

where $\Gamma_k^{(l)} = \Theta_k^{(l)} / \Delta_k^{(l+1)}$. From these two expressions we have

$$\begin{aligned} Y_{k+3}^{(l)} + X_{k+2}^{(l)} + X_{k+1}^{(l)} &= \Gamma_{k+1}^{(l+2)} - \Gamma_k^{(l)} \\ X_{k+1}^{(l+2)} + Y_{k+2}^{(l)} + X_{k+1}^{(l)} &= \Gamma_{k+1}^{(l+2)} - \Gamma_k^{(l)} \end{aligned}$$

and by rearranging, we find a quadrilinear relation in terms of $\Delta_k^{(l)}$ on an 11-point stencil, the HADT equation:

$$\begin{aligned} &\Delta_{k+1}^{(l)} \left(\Delta_k^{(l+4)} \Delta_k^{(l+1)} \Delta_{k-1}^{(l+3)} - \Delta_k^{(l+2)} \Delta_k^{(l+3)} \Delta_{k-1}^{(l+3)} + \Delta_{k-2}^{(l+4)} \Delta_k^{(l+3)} \Delta_{k+1}^{(l+1)} \right) \\ &= \Delta_{k-1}^{(l+4)} \left(\Delta_{k+2}^{(l)} \Delta_k^{(l+1)} \Delta_{k-1}^{(l+3)} - \Delta_k^{(l+2)} \Delta_k^{(l+1)} \Delta_{k+1}^{(l+1)} + \Delta_k^{(l)} \Delta_k^{(l+3)} \Delta_{k+1}^{(l+1)} \right). \end{aligned} \quad (3.18)$$

The HADT equation bears its name as a Higher order Analogue of the Discrete-time Toda equation, due to the similarities between how both equations arise in the theory of formal orthogonal polynomials. We expect the HADT equation to also be integrable and derive a Lax pair for it in the next section. Also note the similarity between the stencils of the two equations (2.8) and (3.18), as displayed in Figure 2.




FIGURE 2. The stencils of the discrete-time Toda equation (2.8) and the HADT equation (3.18)

Next we follow a similar approach as in section 2.1 and derive Lax pairs for the several Hankel identities viewed as integrable systems, using the recurrence relations for the two-variable elliptic orthogonal polynomials. As before the fields will carry two sub-indices; we denote $x = \lambda$, $\Delta_l^{(m)} = \sigma_{l,m}$, $\Theta_l^{(m)} = \rho_{l,m}$, $U_k^{(l)} = u_{k,l}$, etc. By not evaluating the coefficients in the recurrences (3.12) and (3.9), we will find a novel quotient-quotient-difference scheme.

3.5. A Lax pair for the system of lattice equations (3.17). We derive a Lax pair for the system of equations (3.17) which in terms of σ, ρ reads

$$\sigma_{l+1,m-2} \sigma_{l-2,m+2} + \rho_{l,m-2} \sigma_{l-1,m+1} = \sigma_{l,m-2} \sigma_{l-1,m+2} + \sigma_{l,m-1} \rho_{l-1,m}, \quad (3.19a)$$

$$\sigma_{l,m-1} \sigma_{l-1,m+1} + \rho_{l-1,m-1} \sigma_{l,m} = \rho_{l,m-1} \sigma_{l-1,m}. \quad (3.19b)$$

Taking $\Psi_{l,m} = (Q_l^{(m+1)}, P_{l+1}^{(m+1)}, Q_{l+1}^{(m)}, P_{l+2}^{(m)})$ we derive $\Psi_{l+1,m} = L_{l,m}\Psi_{l,m}$, and $\Psi_{l+1,m} = M_{l,m}\Psi_{l,m}$, with

$$L_{l,m} = \begin{pmatrix} \frac{-\sigma_{l+1,m+1}\rho_{l-1,m+1}}{\sigma_{l,m+1}\rho_{l,m+1}} & 1 & 0 & 0 \\ 0 & \frac{-\sigma_{l+2,m}\sigma_{l,m+1}}{\sigma_{l+1,m}\sigma_{l+1,m+1}} & 0 & 1 \\ 0 & 0 & \frac{-\sigma_{l+2,m}\rho_{l,m}}{\sigma_{l+1,m}\rho_{l+1,m}} & 1 \\ 0 & \frac{-\sigma_{l+1,m+2}\sigma_{l,m+1}}{\sigma_{l+3,m}\sigma_{l+1,m}} & 0 & \frac{\sigma_{l+1,m+2}\sigma_{l+1,m+1}-\sigma_{l+1,m}\sigma_{l+1,m+3}}{\sigma_{l,m+3}\sigma_{l+2,m}} \end{pmatrix} \\ + \lambda \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \frac{\sigma_{l+1,m+2}\rho_{l-1,m+1}}{\sigma_{l,m+1}\sigma_{l,m+3}} & 1 + \frac{\rho_{l+1,m+1}\rho_{l,m}}{\sigma_{l,m+3}\sigma_{l+1,m}} & \frac{-\rho_{l+1,m+1}\rho_{l,m}}{\sigma_{l,m+3}\sigma_{l+1,m}} & 0 \end{pmatrix},$$

and

$$M_{l,m} = \begin{pmatrix} \frac{-\rho_{l,m+2}\sigma_{l,m+2}\rho_{l-1,m+1}}{\rho_{l-1,m+2}\sigma_{l,m+1}\sigma_{l,m+3}} & \frac{-\rho_{l,m+2}\rho_{l,m+1}\rho_{l,m}}{\rho_{l-1,m+2}\sigma_{l+1,m}\sigma_{l,m+3}} & \frac{\rho_{l,m+2}\rho_{l,m+1}\rho_{l,m}}{\rho_{l-1,m+2}\sigma_{l+1,m}\sigma_{l,m+3}} & 0 \\ 0 & \frac{\rho_{l+1,m}\sigma_{l,m+1}}{\sigma_{l+1,m}\sigma_{l,m+2}} & \frac{-\sigma_{l+1,m+1}\rho_{l,m}}{\sigma_{l+1,m}\sigma_{l,m+2}} & 0 \\ \frac{-\sigma_{l+1,m+1}\rho_{l-1,m+1}}{\sigma_{l,m+1}\rho_{l,m+1}} & 1 & 0 & 0 \\ 0 & \frac{-\sigma_{l+2,m}\sigma_{l,m+1}}{\sigma_{l+1,m}\sigma_{l+1,m+1}} & 0 & 1 \end{pmatrix} \\ + \lambda^{-1} \begin{pmatrix} 0 & \frac{\sigma_{l,m+1}\rho_{l,m+2}\sigma_{l,m+2}}{\rho_{l-1,m+2}\sigma_{l+1,m}\sigma_{l,m+3}} & \frac{-\rho_{l,m}\sigma_{l,m+2}}{\rho_{l-1,m+2}\sigma_{l+1,m}} & 1 - \frac{\rho_{l,m+2}\sigma_{l-1,m+3}}{\rho_{l-1,m+2}\sigma_{l,m+3}} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

Here we have used the equations (3.19) to reduce the number of terms in the entries of the Lax matrices as much as possible. The compatibility condition $L_{l,m+1}M_{l,m} - M_{l+1,m}L_{l,m} = 0$ is equivalent to the system (3.19). However, to verify that the left hand side vanishes modulo the system, some work has to be done. The equations do not factor out as nicely as in section 2.1.

3.6. A Lax pair for the HADT equation, and the QGD scheme. We first derive a Lax-pair for the following QGD scheme, satisfied by the coefficients ($U_l^{(m)} = u_{l,m}$ etc.) in the recurrence equations (3.9) and (3.12):

$$u_{l+2,m} + v_{l+1,m} + w_{l+1,m+1} = u_{l,m+3} + v_{l+1,m+1} + w_{l+1,m}, \quad (3.20a)$$

$$u_{l,m+3}v_{l,m+1} = v_{l+1,m}u_{l+1,m}, \quad (3.20b)$$

$$u_{l,m+2}w_{l,m} = w_{l+1,m}u_{l+1,m}. \quad (3.20c)$$

Let $\Psi_{l,m} = (P_l^{(m+3)}, P_{l+1}^{(m+2)}, P_{l+2}^{(m+1)}, P_{l+3}^{(m)})$. We use (3.9) and (3.12) to find $\Psi_{l+1,m} = L_{l,m}\Psi_{l,m}$, and $\Psi_{l+1,m} = M_{l,m}\Psi_{l,m}$, where L and M are given

$$L_{l,m} = \begin{pmatrix} -u_{l,m+2} & 1 & 0 & 0 \\ 0 & -u_{l+1,m+1} & 1 & 0 \\ 0 & 0 & -u_{l+2,m} & 1 \\ 0 & 0 & -w_{l+2,m}u_{l+2,m} & -v_{l+2,m} + w_{l+2,m} \end{pmatrix} \\ + \lambda \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ u_{l+1,m+2}u_{l,m+2} & -u_{l+1,m+1} - u_{l+1,m+2} & 1 & 0 \end{pmatrix},$$

and

$$M_{l,m} = \begin{pmatrix} -\frac{u_{l,m+2}}{u_{l,m+3}} & \frac{1}{u_{l,m+3}} & 0 & 0 \\ -u_{l,m+2} & 1 & 0 & 0 \\ 0 & -u_{l+1,m+1} & 1 & 0 \\ 0 & 0 & -u_{l+2,m} & 1 \end{pmatrix} + \lambda^{-1} \begin{pmatrix} 0 & -w_{l,m+1} & 1 + \frac{w_{l+1,m} - v_{l+1,m}}{u_{l,m+3}} & -\frac{1}{u_{l,m+3}} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

Here we have used the scheme (3.20) to simplify the λ^{-1} -part of $M_{l,m}$. The compatibility of these linear systems is equivalent to the QQD scheme (3.20). Note, by substituting the expressions (3.13), (3.10) for u, v, w in terms of σ into the QQD scheme, the first equation is equivalent to the HADT equation, whereas the later two are satisfied identically. Therefore, after the same substitution, the matrices L, M provide a *good* Lax pair for the HADT equation (3.18). The stencils of the Lax matrices $L(\Delta), M(\Delta)$ are depicted in Figures 3, 4.




FIGURE 3.
The stencil of $L_{l,m}$




FIGURE 4.
The stencil of $M_{l,m}$

4. PERIODIC REDUCTION

In [25] a method was given to obtain initial value problems for scalar equations on arbitrary stencils. The question was raised whether a similar construction can be done for systems, and one example was given (the quotient-difference algorithm) where this is the case indeed. Here we will present a systematic approach towards constructing initial value problems for systems of partial difference equations.

Using the method of [25] it is easily shown that the dimension of the \mathbf{s} -periodic initial value problem for the HADT equation is given by the piecewise linear function

$$4\max\{|s_1 + s_2|, |s_1|\}. \quad (4.1)$$

We will show that initial value problems of the same dimension as the initial value problems for the HADT equation, namely (4.1). By constructing In section 4.2 we develop a method to construct initial value problems for systems of lattice equations, which we apply to perform \mathbf{s} -periodic reductions of the intermediate system (3.19). This yields In section 4.3 we present the periodic straight band initial value problems for the QQD scheme, whose well-posedness is proven in Appendix D. Although we have given a method to construct \mathbf{s} -periodic initial value problems for systems, the question remains whether this be done algorithmically. As we will see, one can neither simply determine the dimensionality of the reductions, nor which regions

for \mathbf{s} should be distinguished, from looking at the stencils on which the equations of the system are defined.

4.1. Initial value problems for the HADT equation. To pose initial value problems for equation (1.3) we use the method developed in [25]. By the *S*-polygon of the equation we mean the boundary of the convex hull of the stencil S of the equation (the dotted parallelogram in Figure 5). The *S*-directions are the directions




FIGURE 5. The 11-point stencil of the HADT equation, its convex hull, and two regions in the plane

in the *S*-polygon of the equation: $(1, 0)$ and $(1, -2)$. Therefore, we distinguish two regions in the plane

$$\begin{aligned} R^1 &= \{\mathbf{s} \in \mathbb{Z}^2 : a \geq 0, b > 0, \epsilon = 1 \text{ or } b > 2a, \epsilon = -1\}, \\ R^2 &= \{\mathbf{s} \in \mathbb{Z}^2 : 0 < b < 2a, \epsilon = -1\}, \end{aligned}$$

as in Figure 5. If $\mathbf{s} \in R^1$ or $\mathbf{s} \in R^2$, then performing an \mathbf{s} -periodic reduction yields a well-posed initial value problem, i.e. the \mathbf{s} -periodic solution is obtained by iteration of a single valued mapping. If on the other hand \mathbf{s} is on a boundary between R^1 and R^2 one needs to impose periodicity on the solution, and then one gets a multi-valued mapping, or correspondence. To each of the regions R^i we associate a vector d^i which is a difference between the two points of S , where a line with direction $\mathbf{s} \in R^i$ intersects the *S*-polygon in a single point. For the HADT equation we have

$$d^1 = (0, 4), \quad d^2 = (4, -4). \quad (4.2)$$


Hence the dimension of the \mathbf{s} -periodic initial value problem is, cf. [25, Equation 8],

$$\max_{i=1,2} \left\{ \left| \text{Det} \begin{pmatrix} \mathbf{s} \\ d^i \end{pmatrix} \right| \right\} = 4 \max\{|s_1 + s_2|, |s_1|\}.$$

We give two examples before we present the general \mathbf{s} -periodic initial value problem in terms of \mathbf{s} -reduced variables.

4.1.1. $(2, -1)$ -reduction of the HADT equation. We specify initial values between two parallel lines that squeeze the stencil as in Figure 6. Clearly, given the 8 values x_1, x_2, \dots, x_8 at the black dots one is able to determine the value x_9 at the white dots on the fat-dotted line, using the HADT equation. Thus we get an eight dimensional mapping:

$$\begin{aligned} x_i &\mapsto x_{i+1}, \quad 1 \leq i < 8, \\ x_8 &\mapsto \frac{x_1 x_4 x_7 x_8}{x_2 x_3 x_6} + \frac{x_5 x_7}{x_3} + \frac{x_3 x_8}{x_2} - \frac{x_4 x_5 x_8}{x_2 x_6} - \frac{x_4 x_7^2}{x_3 x_6}. \end{aligned} \quad (4.3)$$


FIGURE 6. $(2,-1)$ -periodic initial values

We can calculate integrals for this mapping by constructing a so called monodromy matrix, which is a product of Lax matrices along a staircase over a one period long distance. Taking the point at the bottom left corner to be the origin $(l, m) = (0, 0)$, we define the *monodromy matrix* to be the inversely ordered product of Lax matrices along a staircase from $(l, m) = (1, 0)$ to $(l, m) = (3, -1)$,

$$\mathcal{L} = (M_{3,-1}^{-1} L_{2,0} L_{1,0}).$$

Using the stencils of the Lax matrices, see Figures 3 and 4, the reader can verify that the matrices in \mathcal{L} depend on initial values only. The characteristic polynomial of \mathcal{L} is

$$\text{Det}(\mu\mathbf{I} - \mathcal{L}) = \mu^2((\mu - \lambda)^2 + 1) + \mu I_1(\mu^2 - \lambda\mu - \lambda) - \lambda(I_2\mu^2 + I_3\mu + 1),$$

where

$$\begin{aligned} I_1 &= \frac{x_3x_6}{x_2x_7} + \frac{x_2x_7}{x_6x_3} - \frac{x_1x_8}{x_6x_3} + \frac{x_1x_4}{x_2x_3} + \frac{x_8x_5}{x_6x_7} - \frac{x_5x_4}{x_2x_7}, \\ I_2 &= \left(\frac{x_5^2}{x_4x_6} + \frac{x_3^2}{x_2x_4} + \frac{x_1x_7}{x_2x_6} - \frac{x_3x_5}{x_2x_6} \right) \left(\frac{x_6^2}{x_5x_7} + \frac{x_4^2}{x_3x_5} + \frac{x_2x_8}{x_3x_7} - \frac{x_4x_6}{x_3x_7} \right), \\ I_3 &= \frac{x_2x_5}{x_3x_4} + \frac{x_6x_3}{x_4x_5} + \frac{x_4x_7}{x_5x_6} + \frac{x_1x_8}{x_3x_6} - \frac{x_2x_7}{x_3x_6}. \end{aligned}$$

Because we subtracted I_1 from the coefficient of $\mu^2\lambda$ in the definition of I_2 it factorizes nicely as JJ' , where J is a 2-integral of the map and J' is the image of J under the map. Remember a k -integral is an integral of the k th power of the map [19]. Indeed, given that $J'' = J$, we find that the product $JJ' = J''J' = (JJ')'$ is an invariant. But now, we can write down another (functionally independent) invariant, namely $J + J'$. Moreover, the integrals I_1, I_2, I_3, I_4 , with

$$I_4 = \frac{x_5^2}{x_4x_6} + \frac{x_3^2}{x_2x_4} + \frac{x_1x_7}{x_2x_6} - \frac{x_3x_5}{x_2x_6} + \frac{x_6^2}{x_5x_7} + \frac{x_4^2}{x_3x_5} + \frac{x_2x_8}{x_3x_7} - \frac{x_4x_6}{x_3x_7},$$

are functionally independent.

4.1.2. *(2,-3)-reduction of the HADT equation.* We specify initial values between two parallel lines that squeeze the stencil as in Figure 7.

Clearly, given the 8 values x_1, x_2, \dots, x_8 at the black dots one is able to determine the value x_9 at the white dots on the fat-dotted line, using the HADT equation. Thus we get an eight dimensional mapping:

$$\begin{aligned} x_i &\mapsto x_{i+1}, \quad 1 \leq i < 8, \\ x_8 &\mapsto \frac{x_6^2}{x_4^2} \left(\frac{x_1x_7}{x_3} - x_5 \right) + \frac{x_7x_5}{x_3}. \end{aligned} \tag{4.4}$$

variables σ_n^p where n runs over a specific range and p runs over the full range $p \in \mathbb{N}_r := \{0, 1, \dots, r-1\}$, cf. [25].

The range of n is determined by the differences associated to each region. For the HADT equation, when $\mathbf{s} \in R^1$ (where $\epsilon = -1$, see Figure 5) the range of n will be from 0 to $n(d^1) - 1 = b \cdot 0 + a \cdot 4 - 1$, and when $\mathbf{s} \in R^2$ we take $0 \leq n < 4(b + \epsilon a) - 1$, using the differences (4.2).

Thus we obtain

Proposition 2. *The HADT equation admits a well-posed \mathbf{s} -periodic initial value problem if (a, b) is not equal to $(1, 0)$ or $(1, -2)$.*

- For $\mathbf{s} \in R^1$ the set $\{\sigma_n^p : n \in \mathbb{N}_{4a}, p \in \mathbb{N}_r\}$ provides a well-posed initial value problem of dimension $4 \mid s_1 \mid$.
- For $\mathbf{s} \in R^2$ the set $\{\sigma_n^p : n \in \mathbb{N}_{4(b+\epsilon a)}, p \in \mathbb{N}_r\}$ provides a well-posed initial value problem of dimension $4 \mid s_1 + s_2 \mid$.

Remark: In section 4.1.1 the initial values $\{\sigma_0^0, \dots, \sigma_7^0\}$ are conveniently denoted $\sigma_n^0 = x_{8-n}$ and the mapping is the backward shift $n \mapsto n-1$, or $(l, m) \mapsto (l-1, m)$. In section 4.1.2 the initial values $\{\sigma_0^0, \dots, \sigma_7^0\}$ are denoted $\sigma_n^0 = x_{n+1}$ and the mapping is the standard one $n \mapsto n+1$, that is $(l, m) \mapsto (l+1, m-1)$.

4.2. Initial value problems for the system of lattice equations (3.19). The stencils of the system (3.19) are depicted in Figure 8. The S -directions in the S -polygons of the equations are $(0, 1)$, $(1, 0)$, $(1, -1)$, and $(1, -2)$. Therefore, we distinguish four regions in the plane, as in Figure 9.




FIGURE 8. The 5-point and 7-point stencils of system (3.19), and their convex hulls




FIGURE 9. Distinguished regions for system (3.19)

For each of the distinguished regions R^i we will project each stencil of the system onto a line with direction $\mathbf{s} \in R^i$. On these lines we will indicate (schematically) the ranges of n -values for both field variables σ, ρ (our ranges start at 0). This procedure is similar to the scalar case, where the difference d^i is equivalent to a range for n . For every equation in a system we need a range (and hence a difference) for every field, and also their relative positions. Moreover, once we have this data, there is still some freedom left how to position the equations with respect to each other. In fact, as we will show in Appendix C, we have to split region R^4 into

two regions where the two equations are positioned differently with respect to each other. Also we will see that for regions R^2 and R^3 we may take the same initial values. However, updating the initial values is done differently in those regions.

For $\mathbf{s} \in R^1$ the ranges are given in Figure 10.




FIGURE 10. \mathbf{s} -Reduction for system (3.19), with $\mathbf{s} \in \mathbb{N}^2$

We now have to find a range of n for each variable σ, ρ such that each equation can be used to update one of the variables. Moreover, we want to minimise the number of field values that have to be calculated to update both fields. Therefore, we move the first equation in Figure 10 over a distance b with respect to the other, see Figure 11.




FIGURE 11. Alternative \mathbf{s} -Reduction for system (3.19), with $\mathbf{s} \in \mathbb{N}^2$

For system (3.19) with $\mathbf{s} \in R^1$ one can see that if we take the range $0 \leq n < 4a + 3b$ for σ and the range $2a + b \leq n < 2a + 2b$ for ρ , then using the equation (3.19a) we can determine the values of ρ at $2a + 2b \leq n \leq 4a + 2b$. Subsequently, the equation (3.19b) can be used to calculate the values of σ at $n = 4a + 3b$. We have obtained the first item in the following proposition, whose complete proof is given in Appendix C.

Proposition 3. *Let $\mathbf{s} \in \mathbb{Z}^2$ be such that (a, b) is not equal to $(1, 0)$ or $(1, -2)$, and let R^i be as in Figure 9. Then system (3.19) admits a well-posed \mathbf{s} -periodic initial value problem.*

- For $\mathbf{s} \in R^1$ the set

$$\{\sigma_n^p, \rho_m^p : n \in \mathbb{N}_{4a+3b}, m - 2a - b \in \mathbb{N}_b, p \in \mathbb{N}_r\}$$

provides a well-posed initial value problem of dimension $4 |s_1 + s_2|$.

- For $\mathbf{s} \in R^2 \cup R^3$ (including the boundary between R^2 and R^3) the set

$$\{\sigma_n^p, \rho_m^p : n \in \mathbb{N}_{4a-b}, m \in \mathbb{N}_b, p \in \mathbb{N}_r\}$$

provides a well-posed initial value problem of dimension $4 |s_1|$.

- For $\mathbf{s} \in R^4$, $b \leq 3a$, the set

$$\{\sigma_n^p, \rho_m^p : n \in \mathbb{N}_{3b-4a}, m + 5a - 2b \in \mathbb{N}_b, p \in \mathbb{N}_r\}$$

provides a well-posed initial value problem of dimension $4 |s_1 + s_2|$.

- For $\mathbf{s} \in R^4$, $b \geq 3a$, the set

$$\{\sigma_n^p, \rho_m^p : n \in \mathbb{N}_{3b-4a}, m+4a-b \in \mathbb{N}_b, p \in \mathbb{N}_r\}$$

provides a well-posed initial value problem of dimension $4|s_1 + s_2|$.

4.2.1. *(2,-1)-reduction of system (3.19)*. We specify initial values between two parallel lines that squeeze the σ -stencil, as in Figure 12. The initial values are denoted $\sigma_n^0 = x_{n+1}$ and $\rho_n^0 = y_{n+1}$. Clearly, from these initial values we can calculate the




FIGURE 12. $(2,-1)$ -periodic initial values

values y_4 and y_2 , y_1 , using equation (3.19b). Then we know enough ρ -values to be able to calculate, using equation (3.19a), x_8 , which (to our surprise) does not depend on y_3 . Therefore, the σ -field is determined by a 7-dimensional mapping, namely

$$\begin{aligned} x_i &\mapsto x_{i+1}, \quad 1 \leq i < 7, \\ x_7 &\mapsto \frac{x_2 x_7}{x_1} - \frac{x_2 x_5 x_6}{x_1 x_4} - \frac{x_3 x_4 x_7}{x_1 x_5} - \frac{x_3^2 x_6^2}{x_1 x_4 x_5}. \end{aligned} \quad (4.6)$$

We know enough values to determine the monodromy matrix $\mathcal{L} = L_0^1 (M_0^1)^{-1} L_0^1$. Its characteristic polynomial is given by

$$\text{Det}(\mu I - \mathcal{L}) = \mu^2((\mu - \lambda)^2 + (\mu - \lambda)I_1 + 1) - \lambda(\mu^2 J J' + \mu I_1 + 1),$$

where

$$\begin{aligned} I_1 &= -\frac{x_3 x_4}{x_1 x_6} + \frac{x_3 x_6}{x_4 x_5} - \frac{x_4 x_5}{x_7 x_2} + \frac{x_1 x_4}{x_2 x_3} + \frac{x_5 x_2}{x_3 x_4} + \frac{x_4 x_7}{x_5 x_6} + \frac{x_5 x_2}{x_1 x_6} \\ &\quad + \frac{x_6 x_3}{x_7 x_2} - \frac{x_5^2 x_2}{x_7 x_1 x_4} - \frac{x_6 x_3^2}{x_7 x_1 x_4} \\ J &= \frac{x_1 x_7 - x_5 x_3}{x_2 x_6} + \frac{x_3^2}{x_2 x_4} + \frac{x_5^2}{x_4 x_6}. \end{aligned}$$

The integrals $I_1, J J', J + J'$ are functionally independent.

4.2.2. *(2,-3)-reduction of system (3.19)*. We specify initial values between two parallel lines that squeeze the σ -stencil, as in Figure 13. From these initial values we can calculate the σ -value (x_6) on the dashed line. And once we have done that we




FIGURE 13. (2,-3)-periodic initial values

can calculate y_4 . We find an 8-dimensional mapping, namely

$$\begin{aligned}
 x_i &\mapsto x_{i+1}, & 1 \leq i < 5, \\
 x_5 &\mapsto x'_5 = \frac{x_4 x_3 + y_1 x_4 - x_3 y_2}{x_1}, \\
 y_i &\mapsto y_{i+1}, & i = 1, 2, \\
 y_3 &\mapsto \frac{x_4 x_5 + y_1 x'_5}{x_3}.
 \end{aligned} \tag{4.7}$$

One has to calculate quite a few other points to be able to evaluate the monodromy matrix, whose characteristic polynomial is given by

$$\text{Det}(\mu I - \mathcal{L}) = \mu^2((\mu - \lambda)^2 + \mu I_1 + \lambda I_2 + 1) - \lambda^3,$$

where the invariants I_1 and I_2 are given in terms of reduced variables in section 5. We have not been able to write any combination of I_1, I_2 in terms of 2-integrals.

4.3. Initial value problems for the QGD scheme. We define regions in the plane as follows

$$\begin{aligned}
 R^1 &= \{\mathbf{s} \in \mathbb{Z}^2 : a \geq 0, b > 0, \epsilon = 1\}, \\
 R^2 &= \{\mathbf{s} \in \mathbb{Z}^2 : 0 < b \leq a, \epsilon = -1\}, \\
 R^3 &= \{\mathbf{s} \in \mathbb{Z}^2 : a \leq b < 2a, \epsilon = -1\}, \\
 R^4 &= \{\mathbf{s} \in \mathbb{Z}^2 : 2a < b \leq 3a, \epsilon = -1\}, \\
 R^5 &= \{\mathbf{s} \in \mathbb{Z}^2 : 3a \leq b, \epsilon = -1\},
 \end{aligned}$$

see Figure 14. We prove Proposition 4 in Appendix D.

Proposition 4. *The QGD scheme (3.20) admits a well-posed \mathbf{s} -periodic initial value problem if (a, b) is not equal to $(1, 0)$ or $(1, -2)$.*

- For $\mathbf{s} \in R^1$ the set

$$\{u_n^p, v_m^p, w_l^p : n \in \mathbb{N}_{3a+2b}, m - 2a \in \mathbb{N}_{a+b}, l - 2a - b \in \mathbb{N}_b, p \in \mathbb{N}_r\}$$

provides a well-posed initial value problem of dimension $4 \lfloor s_1 + s_2 \rfloor$.

- For $\mathbf{s} \in R^2$ the set

$$\{u_n^p, v_m^p, w_l^p : n - b \in \mathbb{N}_{3a-b}, m \in \mathbb{N}_a, l \in \mathbb{N}_b, p \in \mathbb{N}_r\}$$

provides a well-posed initial value problem of dimension $4 \lfloor s_1 \rfloor$.

- For $\mathbf{s} \in R^3$ the set

$$\{u_n^p, v_m^p, w_l^p : n + a - b \in \mathbb{N}_{3a-b}, m \in \mathbb{N}_a, l \in \mathbb{N}_b, p \in \mathbb{N}_r\}$$

provides a well-posed initial value problem of dimension $4 \lfloor s_1 \rfloor$.




FIGURE 14. Distinguished regions for the QQD scheme

- For $\mathbf{s} \in R^4$ the set

$$\{u_n^p, v_m^p, w_l^p : n - 3a + b \in \mathbb{N}_{2b-3a}, m \in \mathbb{N}_{b-a}, l \in \mathbb{N}_b, p \in \mathbb{N}_r\}$$

provides a well-posed initial value problem of dimension $4|s_1 + s_2|$.

- For $\mathbf{s} \in R^5$ the set

$$\{u_n^p, v_m^p, w_l^p : n - 2a \in \mathbb{N}_{2b-3a}, m - a \in \mathbb{N}_{b-a}, l \in \mathbb{N}_b, p \in \mathbb{N}_r\}$$

provides a well-posed initial value problem of dimension $4|s_1 + s_2|$.

In the following reductions we adopt the notation $u_n^0 = x_n, v_n^0 = y_n, w_n^0 = z_n$.

4.3.1. *(2,-1)-reduction of the QQD scheme.* Updating the initial values as described in Appendix D yields the following mapping

$$\begin{aligned} x_i &\mapsto x_{i+1}, & 1 \leq i < 5, \\ x_5 &\mapsto \frac{x_1 y_1}{y_1'}, \\ y_0 &\mapsto y_1, \\ y_1 &\mapsto y_1' = x_1 + y_0 - x_5 - z_0 + \frac{x_5 x_4 z_0}{x_1 x_2}, \\ z_0 &\mapsto \frac{x_4 z_0}{x_1}. \end{aligned} \tag{4.8}$$

The characteristic polynomial of the monodromy matrix is given by

$$\text{Det}(\mu I - \mathcal{L}) = \mu^2((\mu - \lambda)^2 + (\mu - \lambda)I_1 - \lambda J J' + I_2) + \lambda(\mu I_3 - I_4),$$

where

$$\begin{aligned}
I_1 &= (y_0 - z_0 + x_1) \frac{z_0 x_4}{x_1} + y_1 (z_0 + x_5 - y_0), \\
I_2 &= z_0 x_4 x_5 y_1, \\
I_3 &= (z_0 x_2 - x_5 y_1 - x_2 x_1 - y_0 x_2) x_3 x_4 - x_1 y_1 (x_5 x_4 + x_2 x_5 + x_2 x_3), \\
I_4 &= x_1 x_2 x_3 x_4 x_5 y_1, \\
J &= x_1 + x_3 + y_0 - z_0.
\end{aligned}$$

The integrals $I_1, I_2, I_3, I_4, I_5 = JJ', I_6 = J + J'$ are functionally independent.

4.3.2. *(2,-3)-reduction of the QQD scheme.* Updating the initial values as described in Appendix D yields the following mapping

$$\begin{aligned}
x_i &\mapsto x_{i+1}, \quad i = 1, 2, \\
x_3 &\mapsto x'_3 = \frac{y_1 x_1}{y_0}, \\
y_0 &\mapsto y_1 \\
y_1 &\mapsto y_0 + z_2 - z_0, \\
z_i &\mapsto z_{i+1}, \quad i = 0, 1, \\
z_2 &\mapsto \frac{x'_3 z_0}{x_3}.
\end{aligned} \tag{4.9}$$

The characteristic polynomial of the monodromy matrix is given by

$$\begin{aligned}
\text{Det}(\mu I - \mathcal{L}) &= \mu^2((\mu - \lambda)^2 + \mu(I_4 - I_3 + (I_5 - I_2 + 1)I_1) + \lambda(I_3 + (2I_2 - 3)I_1)) \\
&\quad + \lambda^2(\mu(3 - I_2) - \lambda)I_1,
\end{aligned}$$

where

$$\begin{aligned}
I_1 &= \frac{x_2 x_1 x_3}{y_0}, \\
I_2 &= \frac{z_2}{x_3} + \frac{y_0}{x_3} + \frac{y_0}{x_1} + \frac{y_1}{x_2}, \\
I_3 &= x_2 z_0 - x_2 y_0 - x_2 z_2 + z_1 x_3 - x_3 y_1 - y_1 x_1, \\
I_4 &= y_1 z_0 - y_1 y_0 + y_0 z_1 - z_1 z_0, \\
I_5 &= \frac{z_0 z_1 z_2}{x_3}.
\end{aligned}$$

5. DIMENSIONAL REDUCTION

In this section we present the explicit formulae for the reduced variables, required to implement the dimensional reduction of the mappings obtained in section 4 from periodic reduction from the HADT equation and from the system (3.19). These formulae arise from the symmetries of the partial difference equations. In certain cases an extra scaling symmetry of the integrals has to be used to obtain the dimensional reduction.

5.1. **The HADT equation.** Equation (3.18) is invariant under the following continuous symmetries: $u \mapsto u\epsilon$ and $u \mapsto u\epsilon^{(-1)^l}$. The first yields a symmetry of

the mappings, whereas the second gives us a 2-symmetry of the mappings. Their generators are

$$\sum_{i=1}^8 x_i \frac{\partial}{\partial x_i}, \quad \sum_{i=1}^8 (-1)^i x_i \frac{\partial}{\partial x_i}.$$

A set of joint invariants of these vector fields is given by

$$\left\{ w_1 = \frac{x_1 x_4}{x_2 x_3}, w_2 = \frac{x_2 x_5}{x_3 x_4}, w_3 = \frac{x_3 x_6}{x_4 x_5}, w_4 = \frac{x_4 x_7}{x_5 x_6}, w_5 = \frac{x_5 x_8}{x_6 x_7}, w_6 = \frac{x_6 x_1}{x_7 x_8} \right\}.$$

In these variables the $\mathbf{s} = (2, -1)$ mapping (4.3) reads

$$\begin{aligned} w_i &\mapsto w_{i+1}, \quad 1 \leq i < 5, \\ w_5 &\mapsto \frac{f}{w_2 w_3 w_4 w_5}, \quad f = w_2 - w_5 + (w_3 - w_2)w_4 + w_1 w_2 w_3 w_4 w_5, \\ w_6 &\mapsto \frac{w_2 w_4 w_6}{w_1 f}. \end{aligned} \tag{5.1}$$

which has three functionally independent invariants

$$I_1 = w_1 + w_5 + w_2 w_4 (1 - w_1 w_3 w_5) + \frac{w_3 - 1}{w_2 w_3 w_4}, \tag{5.2a}$$

$$I_2 = \frac{(w_1 w_2 w_3 w_4 + w_2 + w_3 - 1)(w_2 w_3 w_4 w_5 + w_3 + w_4 - 1)}{w_2 w_3 w_4}, \tag{5.2b}$$

$$I_3 = w_2 + w_3 + w_4 - w_4 w_2 + w_1 w_2 w_3 w_4 w_5. \tag{5.2c}$$

The 2-integral J is scaled by the generator of the second symmetry and so it does not survive the reduction. The $\mathbf{s} = (2, -3)$ mapping (4.4) reduces to the six dimensional mapping

$$\begin{aligned} w_i &\mapsto w_{i+1}, \quad 1 \leq i < 5, \\ w_5 &\mapsto \frac{g}{w_4 w_5}, \quad g = w_1 w_2 w_3 + \frac{1}{w_3} - \frac{1}{w_4} \\ w_6 &\mapsto \frac{w_4 w_6}{w_1 w_3 g} \end{aligned} \tag{5.3}$$

and has three functionally independent invariants

$$I_1 = w_3(w_1 w_2 + w_2 w_4 + w_4 w_5) + \frac{1}{w_3},$$

$$I_2 = w_2 w_3^2 w_4 (w_1 w_2 + w_1 w_5 + w_4 w_5) - w_2 w_3 - w_3 w_4 + w_2 w_4,$$

$$JJ' = w_3(w_1 w_2 w_3 w_4 - 1)(w_2 w_3 w_4 w_5 - 1).$$

5.2. The lattice system (3.19). For the case $\mathbf{s} = (2, -1)$ the mapping has a symmetry and a 2-symmetry, which arise from scaling symmetries of the system. Their generators are given by

$$\sum_{i=1}^7 x_i \frac{\partial}{\partial x_i}, \quad \sum_{i=1}^7 (-1)^i x_i \frac{\partial}{\partial x_i}.$$

The integrals admit an extra scaling, which is neither a symmetry, nor a k -symmetry, of the mapping, namely

$$x_1 \frac{\partial}{\partial x_1} + x_2 \frac{\partial}{\partial x_2} - x_5 \frac{\partial}{\partial x_5} - x_6 \frac{\partial}{\partial x_6} - 2x_7 \frac{\partial}{\partial x_7}.$$

Still, we can use joint invariants of these three vector fields,

$$w_1 = \frac{x_1x_4}{x_2x_3}, w_2 = \frac{x_5x_2}{x_3x_4}, w_3 = \frac{x_3x_6}{x_5x_4}, w_4 = \frac{x_4x_7}{x_5x_6},$$

to reduce the dimension of the mapping by 3 dimensions. This is due to the fact that if we reduce the third vector field by the first two the resulting vector field is a symmetry of the reduced mapping. The mapping (4.6) reduces to a four dimensional mapping

$$\begin{aligned} w_i &\mapsto w_{i+1}, \quad 1 \leq i < 4, \\ w_4 &\mapsto \frac{w_2w_4 - w_2 - w_3 - w_4}{w_1w_2w_3w_4}, \end{aligned} \tag{5.4}$$

which admits two functionally independent integrals

$$\begin{aligned} I_1 &= w_1 + w_2 + w_3 + w_4 + \frac{w_1w_3 + w_2w_4 - w_1 - w_2 - w_3 - w_4}{w_1w_2w_3w_4}, \\ JJ' &= (w_1 + w_2 + w_3 + w_4 - w_1w_3 - w_1w_4 - w_2w_4) \frac{w_1w_2w_3w_4 + w_2 + w_3 - 1}{w_1w_2w_3w_4}. \end{aligned}$$

Similarly we find three reductions for the $\mathfrak{s} = (2, -3)$ case. Here choosing variables

$$w_1 = \frac{y_1}{x_3}, w_2 = \frac{y_2}{x_4}, w_3 = \frac{y_3}{x_5}, w_4 = \frac{x_2x_3}{x_1x_4}, w_5 = \frac{x_3x_4}{x_2x_5}$$

reduces the mapping (4.7) to the five dimensional mapping

$$\begin{aligned} w_i &\mapsto w_{i+1}, \quad i = 1, 2, 4, \\ w_3 &\mapsto w_1 + w'_5, \\ w_5 &\mapsto w'_5 = \frac{1}{(1 + w_1 - w_2)w_4w_5}. \end{aligned} \tag{5.5}$$

This mapping admits the two integrals

$$\begin{aligned} I_1 &= w_1(w_2 - w_1) + w_2(w_3 - w_2) + w_3(w_1 - w_3) \\ &\quad + w_4(w_2 - w_1 - 1) + w_5(w_3 - w_2 - 1) - \frac{1}{w_4w_5}, \\ I_2 &= ((w_3 - w_2 + w_4)(w_5 - w_3 + w_1) - w_5)(w_1 - w_2) \\ &\quad + \frac{w_3 - w_2 + w_4}{w_4w_5} + w_4(w_5 - w_3 + w_1). \end{aligned}$$

APPENDIX A. SOME DETERMINANTAL IDENTITIES

In the establishment of the recursive structure we need a number of determinantal identities, which we derive using the Sylvester Identity. So we present a proof of the Sylvester identity, which was first presented by Kowalewski [28], Bareiss [4] and Malaschonok [32, 33] and these seven proofs are presented together in [2].

We consider an $(n+m) \times (n+m)$ matrix R with elements r_{ij} and determinant $|R|$, also written $\text{Det}(R)$. Then we partition R and factor by block triangularization such that

$$R = \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} A & 0 \\ C & \mathbf{1} \end{pmatrix} \cdot \begin{pmatrix} \mathbf{1} & A^{-1}B \\ 0 & D - CA^{-1}B \end{pmatrix} \quad (\text{A.1})$$

where A is a nonsingular square matrix of order n , then

$$|R| = |A| \cdot |D - CA^{-1}B|. \quad (\text{A.2})$$

If we multiply both sides by $|A|^{m-1}$, this becomes

$$|A|^{m-1}|R| = ||A|(D - CA^{-1}B)|$$

because the determinant on the right side of (A.2) is of order m . We can reduce this equation further to

$$|A|^{m-1}|R| = ||A|D - C\tilde{A}B|, \quad (\text{A.3})$$

since $A^{-1} = \frac{\tilde{A}}{|A|}$ (where \tilde{A} represents the adjugate matrix of the inverse matrix A^{-1}), and the determinant of A is assumed to be $\neq 0$. Specifying some entries in (A.1), taking A to be an $n \times n$ block and D to be an $m \times m$ block, we have the formula:

$$(\text{Det}(A))^{m-1} \begin{vmatrix} A & | & b_1 & \dots & b_m \\ - & + & - & - & - \\ c_1^t & | & & & \\ \vdots & | & & D & \\ c_m^t & | & & & \end{vmatrix} = \text{Det}_{m \times m} \left\{ \text{Det}_{n \times n}(A) D_{ij} - (c_i^t \tilde{A} b_j) \right\}_{i,j=1,\dots,m} \quad (\text{A.4})$$

in which the full matrix is supplemented with m n -component column vectors b_i and m n -component row-vectors c_i^t . If we consider the case $m = 2$ ie. the removal of two rows and columns, then we get then determinant identity

$$\begin{aligned} (\text{Det}(A)) \begin{vmatrix} A & | & b_1 & b_2 \\ - & + & - & - \\ c_1^t & | & d_{11} & d_{12} \\ c_2^t & | & d_{21} & d_{22} \end{vmatrix} &= \text{Det}_{2 \times 2} \left\{ (\text{Det}(A)) \begin{pmatrix} d_{11} & d_{12} \\ d_{21} & d_{22} \end{pmatrix} - \begin{pmatrix} c_1^t \tilde{A} b_1 & c_1^t \tilde{A} b_2 \\ c_2^t \tilde{A} b_1 & c_2^t \tilde{A} b_2 \end{pmatrix} \right\} \\ &= [\text{Det}(A) d_{11} - c_1^t \tilde{A} b_1] [\text{Det}(A) d_{22} - c_2^t \tilde{A} b_2] \\ &\quad - [\text{Det}(A) d_{21} - c_2^t \tilde{A} b_1] [\text{Det}(A) d_{12} - c_1^t \tilde{A} b_2], \end{aligned}$$

which can be symbolically written as:

$$\begin{array}{c}
 \left| \begin{array}{c} \\ \\ \end{array} \right| \times \left| \begin{array}{c} \\ \\ \end{array} \right| = \left| \begin{array}{c} \\ \\ \end{array} \right| \times \left| \begin{array}{c} \\ \\ \end{array} \right| \\
 - \left| \begin{array}{c} \\ \\ \end{array} \right| \times \left| \begin{array}{c} \\ \\ \end{array} \right|
 \end{array} \tag{A.5}$$

(where the red lines denote rows and columns omitted from the original determinant). It is then necessary to reorder the position of the row and column to tailor the identity to our requirements.

While (A.5) is the key identity by which the recurrence structure for ordinary one-variable orthogonal polynomials is obtained, for the elliptic two-variable orthogonal polynomials we need (in addition to (A.5)), determinantal identities involving the simultaneous removal of more than two rows and columns. Thus, the main identity we use from the general formula (A.4) will be the case $m = 3$, leading to the different recurrence relations for (3.3) and (3.7). So considering $m = 3$ we obtain from (A.4) the following 3-row/column Sylvester type identity:

$$\begin{array}{c}
 \left| \begin{array}{c} \\ \\ \end{array} \right| \times \left| \begin{array}{c} \\ \\ \end{array} \right| = \left| \begin{array}{c} \\ \\ \end{array} \right| \times \left| \begin{array}{c} \\ \\ \end{array} \right| \\
 - \left| \begin{array}{c} \\ \\ \end{array} \right| \times \left| \begin{array}{c} \\ \\ \end{array} \right| + \left| \begin{array}{c} \\ \\ \end{array} \right| \times \left| \begin{array}{c} \\ \\ \end{array} \right|,
 \end{array} \tag{A.6}$$

which is obtained from the expansion of (A.4) for $m = 3$ after recombination of terms using the earlier 2-row/column Sylvester identity (A.5).

APPENDIX B. INTERMEDIATE DETERMINANTS

Previously we have chosen to apply a 3 row/column Sylvester identity to the $P_k^{(l)}$ and $Q_k^{(l)}$ polynomials since this identity (A.6) does not introduce new determinants. However for some 2 row/column Sylvester identities it is possible to control new determinants, which can be removed to still give equations in term $P_k^{(l)}$ and $Q_k^{(l)}$ only. These “*intermediate*” determinants are introduced here.

B.1. The linear $P_k^{(l)}$ equation. By considering the $P_k^{(l)}$ with the first column and penultimate row removed we get the *intermediate* quantity:

$$T_{k-1}^{(l)}(x, y) \equiv \left| \begin{array}{cccccc} \langle \mathbf{e}_l, \mathbf{e}_2 \rangle & \langle \mathbf{e}_l, \mathbf{e}_3 \rangle & \cdots & \cdots & \langle \mathbf{e}_l, \mathbf{e}_k \rangle \\ \langle \mathbf{e}_{l+1}, \mathbf{e}_2 \rangle & \langle \mathbf{e}_{l+1}, \mathbf{e}_3 \rangle & \cdots & \cdots & \langle \mathbf{e}_{l+1}, \mathbf{e}_k \rangle \\ \vdots & \vdots & & & \vdots \\ \vdots & \vdots & & & \vdots \\ \langle \mathbf{e}_{l+k-3}, \mathbf{e}_2 \rangle & \langle \mathbf{e}_{l+k-3}, \mathbf{e}_3 \rangle & \cdots & \cdots & \langle \mathbf{e}_{l+k-3}, \mathbf{e}_k \rangle \\ \mathbf{e}_2 & \mathbf{e}_3 & \cdots & \cdots & \mathbf{e}_k \end{array} \right| / \Pi_{k-2}^{(l)}, \quad (\text{B.1})$$

together with a corresponding Hankel determinant:

$$\Pi_{k-1}^{(l)} \equiv \left| \begin{array}{cccccc} \langle \mathbf{e}_l, \mathbf{e}_2 \rangle & \langle \mathbf{e}_l, \mathbf{e}_3 \rangle & \cdots & \cdots & \langle \mathbf{e}_l, \mathbf{e}_k \rangle \\ \langle \mathbf{e}_{l+1}, \mathbf{e}_2 \rangle & \langle \mathbf{e}_{l+1}, \mathbf{e}_3 \rangle & \cdots & \cdots & \langle \mathbf{e}_{l+1}, \mathbf{e}_k \rangle \\ \vdots & \vdots & & & \vdots \\ \vdots & \vdots & & & \vdots \\ \langle \mathbf{e}_{l+k-2}, \mathbf{e}_2 \rangle & \langle \mathbf{e}_{l+k-2}, \mathbf{e}_3 \rangle & \cdots & \cdots & \langle \mathbf{e}_{l+k-2}, \mathbf{e}_k \rangle \end{array} \right|. \quad (\text{B.2})$$

Using the usual Sylvester identity we can now derive the following two equations

$$\left| \begin{array}{c} \boxed{P_k^{(l)}} \end{array} \right| \Rightarrow P_k^{(l)} = T_{k-1}^{(l+1)} - \frac{\Delta_{k-2}^{(l+1)} \Pi_{k-1}^{(l)}}{\Delta_{k-1}^{(l)} \Pi_{k-2}^{(l+1)}} P_{k-1}^{(l+1)}, \quad (\text{B.3a})$$

$$\left| \begin{array}{c} \boxed{P_k^{(l)}} \end{array} \right| \Rightarrow P_k^{(l)} = T_{k-1}^{(l)} - \frac{\Delta_{k-2}^{(l)} \Pi_{k-1}^{(l)}}{\Delta_{k-1}^{(l)} \Pi_{k-2}^{(l)}} P_{k-1}^{(l)}. \quad (\text{B.3b})$$

Eliminating the $T_k^{(l)}$ polynomials in favor of the $P_k^{(l)}$ polynomials and using a Hankel identity

$$\left| \begin{array}{c} \boxed{\Pi_k^{(l)}} \end{array} \right| \Rightarrow \Pi_k^{(l)} \Delta_{k-2}^{(l+3)} = \Pi_{k-1}^{(l)} \Delta_{k-1}^{(l+3)} - \Pi_{k-1}^{(l+1)} \Delta_{k-1}^{(l+2)} \quad (\text{B.4})$$

allows the derivation of the linear relation in $P_k^{(l)}$ (3.12).

B.2. The bilinear relation in $\Delta_k^{(l)}, \Theta_k^{(l)}$. It is possible to combine some three-term bilinear Hankel relations, to give a three-term relation in terms of $\Delta_k^{(l)}, \Theta_k^{(l)}$. This particular identity is achieved with the introduction of a new *intermediate* determinant Ξ which is essentially Θ with the first column and last row removed.

$$\left| \begin{array}{|c|} \hline \hline \Delta_k^{(l)} \\ \hline \hline \end{array} \right| \Rightarrow \Delta_k^{(l)} \Pi_{k-2}^{(l+2)} = \Theta_{k-1}^{(l)} \Pi_{k-1}^{(l+1)} - \Delta_{k-1}^{(l+1)} \Xi_{k-1}^{(l)} \quad (\text{B.5a})$$

$$\left| \begin{array}{|c|} \hline \hline \Theta_k^{(l)} \\ \hline \hline \end{array} \right| \Rightarrow \Theta_k^{(l)} \Pi_{k-2}^{(l+2)} = \Theta_{k-1}^{(l)} \Pi_{k-1}^{(l+2)} - \Delta_{k-1}^{(l+2)} \Xi_{k-1}^{(l)} \quad (\text{B.5b})$$

where

$$\Xi_{k-1}^{(l)} \equiv \begin{vmatrix} \langle \mathbf{e}_l, \mathbf{e}_2 \rangle & \langle \mathbf{e}_l, \mathbf{e}_3 \rangle & \cdots & \cdots & \langle \mathbf{e}_l, \mathbf{e}_k \rangle \\ \langle \mathbf{e}_{l+2}, \mathbf{e}_2 \rangle & \langle \mathbf{e}_{l+2}, \mathbf{e}_3 \rangle & \cdots & \cdots & \langle \mathbf{e}_{l+2}, \mathbf{e}_k \rangle \\ \vdots & \vdots & & & \vdots \\ \vdots & \vdots & & & \vdots \\ \langle \mathbf{e}_{l+k-1}, \mathbf{e}_2 \rangle & \langle \mathbf{e}_{l+k-1}, \mathbf{e}_3 \rangle & \cdots & \cdots & \langle \mathbf{e}_{l+k-1}, \mathbf{e}_k \rangle \end{vmatrix}. \quad (\text{B.6})$$

Ξ is then eliminated in the combination of the two bilinear relations (B.5).

$$\Delta_{k-1}^{(l+1)} (\Theta_{k-1}^{(l)} \Pi_{k-1}^{(l+2)} - \Theta_k^{(l)} \Pi_{k-2}^{(l+2)}) = \Delta_{k-1}^{(l+2)} (\Theta_{k-1}^{(l)} \Pi_{k-1}^{(l+1)} - \Delta_k^{(l)} \Pi_{k-2}^{(l+2)}) \quad (\text{B.7})$$

This equation can be expanded (in the third term) using a third bilinear equation

$$\left| \begin{array}{|c|} \hline \hline \Delta_k^{(l)} \\ \hline \hline \end{array} \right| \Rightarrow \Delta_k^{(l)} \Pi_{k-2}^{(l+1)} = \Delta_{k-1}^{(l)} \Pi_{k-1}^{(l+1)} - \Delta_{k-1}^{(l+1)} \Pi_{k-1}^{(l)}. \quad (\text{B.8})$$

Thus we are left with an equation in terms of Δ and Θ only.

$$\begin{aligned} \Delta_{k-1}^{(l+1)} (\Theta_{k-1}^{(l)} \Pi_{k-1}^{(l+2)} - \Theta_k^{(l)} \Pi_{k-2}^{(l+2)}) &= \Theta_{k-1}^{(l)} (\Delta_{k-1}^{(l+1)} \Pi_{k-1}^{(l+2)} - \Delta_k^{(l+1)} \Pi_{k-2}^{(l+2)}) \\ &\quad - \Delta_{k-1}^{(l+2)} \Delta_k^{(l)} \Pi_{k-2}^{(l+2)} \end{aligned}$$

After we have canceled the necessary terms, we are left with (3.17b).

B.3. Relations in $P_k^{(l)}$ and $Q_k^{(l)}$. For relations involving the Hankel determinants $P_k^{(l)}$ and $Q_k^{(l)}$ we can derive a recurrence relation $xQ_k^{(l)}$ and two linear relations. By

implementing a cutting of rows and columns on the matrix for $P_k^{(l)}$ according to:

and applying the 3-row/column Sylvester identity in that situation, we obtain:

$$P_k^{(l)} = xQ_{k-2}^{(l+2)} - \frac{\Delta_{k-2}^{(l)}\Theta_{k-2}^{(l+2)}}{\Delta_{k-1}^{(l)}\Theta_{k-3}^{(l+2)}} P_{k-1}^{(l)} + \frac{\Theta_{k-2}^{(l)}\Delta_{k-2}^{(l+2)}}{\Delta_{k-1}^{(l)}\Theta_{k-3}^{(l+2)}} Q_{k-1}^{(l)} \quad , \quad l \neq 0, 1 . \quad (\text{B.9})$$

For the linear relations we first obtain

which by eliminating the $T_k^{(l)}$ using (B.3a) together with the Hankel identity (B.5a) leads to

$$Q_k^{(l)} = P_k^{(l+1)} + \frac{\Delta_k^{(l)}\Delta_{k-2}^{(l+2)}}{\Theta_{k-1}^{(l)}\Delta_{k-1}^{(l+1)}} P_{k-1}^{(l+2)} . \quad (\text{B.11})$$

This three term equation in terms of $Q_k^{(l)}$ and $P_k^{(l)}$ is similar to (3.12). To find another of this type of equation we must first introduce the *intermediate* polynomials $S_{k-1}^{(l)}$:

$$S_{k-1}^{(l)}(x, y) \equiv \left| \begin{array}{cccccc} \langle \mathbf{e}_l, \mathbf{e}_2 \rangle & \langle \mathbf{e}_l, \mathbf{e}_3 \rangle & \cdots & \cdots & \langle \mathbf{e}_l, \mathbf{e}_k \rangle \\ \langle \mathbf{e}_{l+2}, \mathbf{e}_2 \rangle & \langle \mathbf{e}_{l+2}, \mathbf{e}_3 \rangle & \cdots & \cdots & \langle \mathbf{e}_{l+2}, \mathbf{e}_k \rangle \\ \vdots & \vdots & & & \vdots \\ \vdots & \vdots & & & \vdots \\ \langle \mathbf{e}_{l+k-2}, \mathbf{e}_2 \rangle & \langle \mathbf{e}_{l+k-2}, \mathbf{e}_3 \rangle & \cdots & \cdots & \langle \mathbf{e}_{l+k-2}, \mathbf{e}_k \rangle \\ \mathbf{e}_2 & \mathbf{e}_3 & \cdots & \cdots & \mathbf{e}_k \end{array} \right| / \Gamma_{k-2}^{(l)} . \quad (\text{B.12})$$

Then applying the following Sylvester identities to $P_k^{(l)}$ and $Q_k^{(l)}$, we subsequently obtain the relations:

$$\left| \begin{array}{c|c|c} & & \\ \hline & Q_k^{(l)} & \\ \hline & & \\ \hline & & \end{array} \right| \Rightarrow Q_k^{(l)} = S_{k-1}^{(l)} - \frac{\Gamma_{k-1}^{(l)} \Theta_{k-2}^{(l)}}{\Gamma_{k-2}^{(l)} \Theta_{k-1}^{(l)}} Q_{k-1}^{(l)}, \quad (\text{B.13a})$$

$$\left| \begin{array}{c|c|c} & & \\ \hline & P_k^{(l)} & \\ \hline & & \\ \hline & & \end{array} \right| \Rightarrow P_k^{(l)} = S_{k-1}^{(l)} - \frac{\Pi_{k-1}^{(l)} \Theta_{k-2}^{(l)}}{\Gamma_{k-2}^{(l)} \Delta_{k-1}^{(l)}} Q_{k-1}^{(l)}. \quad (\text{B.13b})$$

We eliminate $S_{k-1}^{(l)}$ and making use of the Hankel determinant identity

$$\left| \begin{array}{c|c|c} & & \\ \hline & \Delta_k^{(l)} & \\ \hline & & \\ \hline & & \end{array} \right| \Rightarrow \Delta_k^{(l)} \Xi_{k-2}^{(l)} = \Delta_{k-1}^{(l)} \Xi_{k-1}^{(l)} - \Theta_{k-1}^{(l)} \Pi_{k-1}^{(l)} \quad (\text{B.14})$$

which leaves:

$$Q_k^{(l)} = P_k^{(l)} - \frac{\Delta_k^{(l)} \Theta_{k-2}^{(l)}}{\Delta_{k-1}^{(l)} \Theta_{k-1}^{(l)}} Q_{k-1}^{(l)}. \quad (\text{B.15})$$

APPENDIX C. s -PERIODIC INITIAL VALUE PROBLEMS FOR SYSTEM (3.19)

Proposition 3 is obtained by considering the ranges of the n -values for the equations with periods taken in the remaining regions. When $\mathbf{s} \in R^2$ (see Figure 9) we get the following reduction.




FIGURE 15. s -Reduction for system (3.19), with $\mathbf{s} \in R^2$

So we first need to calculate the values of ρ for $b \leq n \leq 2a - b$. This can be done using equation (3.17a) because the values of σ at n smaller than $2a + (2a - b - b) = 4a - 2b$ are given initially. Then the values of σ at $n = 4a - b$ are determined by (3.17b). When $\mathbf{s} \in R^3$ we get




FIGURE 16. s -Reduction for the system (3.19), with $\mathbf{s} \in R^3$

In Figure 16 the dashed line indicates that $a + b$ can either be to the left, or to the right of $4a - b$. We have $a + b < 4a - b$ in the region $3a > 2b > 2a$. Here it does not matter whether one first calculates the values of ρ at $n = b$, using equation (3.17a), or the values of σ at $n = 4a - b$, using equation (3.17b). In the region $3a \leq 2b < 4a$ we have $a + b \geq 4a - b$. Here we need first calculate the values of σ for $4a - b \leq n \leq a + b$, using equation (3.17b). This can be done indeed, because the values of ρ with $n \leq 2a - b + (a + b) - (4a - b) = b - a$ are given initially. Subsequently the values of ρ at $n = b$ can be obtained.

When $\mathbf{s} \in R^4$ we also need to distinguish two cases, but here this leads to different initial value problems. When $2a > b \leq 3a$ we can first calculate the values of σ at $n = 3b - 4a$, and then the values of ρ at $n = 3b - 5a$, see Figure 17.




FIGURE 17. s -Reduction for the system (3.19), with $\mathbf{s} \in R^4$, $b \leq 3a$

When $b \geq 3a$ we need first calculate the values of ρ at $n = 2b - 4a$, and then the values of σ at $n = 3b - 4a$, see Figure 18.




FIGURE 18. s -Reduction for the system (3.19), with $s \in \mathbb{R}^4$, $b \geq 3a$

We leave it to the reader to check that the given initial value problems can also be updated in the negative n -direction. In Appendix E we illustrate with an example that this is not necessarily the case, and thus has to be verified separately.

APPENDIX D. \mathbf{s} -PERIODIC INITIAL VALUE PROBLEMS FOR THE QQD SCHEME

The stencils of the QQD scheme (3.20) are depicted in Figure 19.




FIGURE 19. Graphical representation of the QQD scheme

To describe initial value problems for the QQD scheme we follow the following procedure. We project the stencils of the three equations onto lines with directions in different regions and translate them in such a way that, given the values of the fields for certain ranges of n , each equation determines a value for an update of one of the fields. The proof of Proposition consists of 5 pictures, one for each region. For every picture one has to check the following:

- (1) that the ranges of n -values for u, v, w correspond to each equation in the system, for any chosen direction \mathbf{s} in the particular region, this includes checking the order of the linear expressions in a, b .
- (2) that, given the initial values for u, v, w , every equation can be used to update one of the fields (in both directions).




FIGURE 20. \mathbf{s} -Reduction of the QQD scheme, $\mathbf{s} \in R^1$

We will perform check (1) and check (2, positive direction) for the first picture, and check (2, positive direction) for the other pictures. This enables one to perform \mathbf{s} -reduction for any given $\mathbf{s} \in \cup_i R^i$. The rest of the proof is left to the reader.

When $\epsilon = 1$, as in region R^1 , with one step to the right (downwards) n increases by b (by a). Let n be equal to 0 on a line with direction $\mathbf{s} \in R^1$ through the first point u in the first equation of the system (the upper-left point in Figure 19). We let the upper-left point from the second equation coincide with the one of the first equation and we let the upper-left point from the third equation be one step to the right, so that it is at distance b from the others, as in Figure 21. Moving the line over the Figure 21 in downward direction, depending on the direction \mathbf{s} , the line will either first cross this point at distance b , or the point in the second equation at distance $2a$. This is indicated in Figure 20 by the dashed line between b and $2a$. Next the line moves over the point with distance $2a + b$, and then it depends on \mathbf{s} whether $2a + 2b$ or $3a + b$ is encountered first. This is again indicated by a dashed




FIGURE 21. The equations of the QQD scheme on top of each other, and a line with direction $(2, 1) \in R^1$.

line. Thus, Figure 20 represents the ranges of the n -values of the three fields of the three equations in the QQD scheme and their relative positions, for $\mathbf{s} \in R^1$.

Next, suppose initially we are given the values of u at $0 \leq n < 3a + 2b$, of v at $2a \leq n < 3a + b$, and of w at $2a + b \leq n < 2a + 2b$. Then, for all $\mathbf{s} \in R^1$, we can use the second equation to determine the value of v at $n = 3a + b$. If $a < b$, then we know all values of w at $2a + b \leq n < 3a + b$, and hence we can use the first equation to calculate the value of u at $n = 3a + 2b$. Then, we can calculate the value of w at $n = 2a + 2b$. However, if $a \geq b$, then we first have to use the last equation to calculate all values of w at $2a + 2b \leq n \leq 3a + b$, before we can use the first equation to calculate the value of u at $n = 3a + 2b$. In both cases, the initial value problem is well-posed.




FIGURE 22. \mathbf{s} -Reduction of the QQD scheme, $\mathbf{s} \in R^2$

In R^2 we first use the third equation to determine the values of w at $b \leq n \leq a$, then the first equation to determine the values of v at $n = a$, and finally the second equation to determine the values of u at $n = 3a$.




FIGURE 23. \mathbf{s} -Reduction of the QQD scheme, $\mathbf{s} \in R^3$

In R^3 the values of u at $n = 2a$ and the values of v at $n = a$ can be determined independently, using the second and first equation, respectively. After having determined the values of u at $n = 2a$ we are able to find the w at $n = b$, using the third equation.




FIGURE 24. \mathfrak{s} -Reduction of the QQD scheme, $\mathfrak{s} \in R^4$

In R^4 the values of u at $n = b$ and the values of v at $n = b - a$ can be determined independently, using the first and second equation, respectively. After having determined the values of u at $n = b$ we are able to find the w at $n = b$, using the third equation.




FIGURE 25. \mathfrak{s} -Reduction of the QQD scheme, $\mathfrak{s} \in R^5$

In R^5 the values of v and w at $n = b$ can be determined independently, using the second and third equation, respectively. Next the values of u at $n = 2b - a$ can be determined using the first equation.

APPENDIX E. A NON-INVERTIBLE REDUCTION OF THE QQD SCHEME

Using the same notation as in section 4.3, consider initial values $\{x_0, x_1, x_2, x_3, x_4, y_0, y_1, z_0, z_1\}$ for the $(2, -1)$ reduction of the QQD scheme. We use equation (3.20b) to determine x_5 , equation (3.20c) to determine z_2 and equation (3.20a) to determine y_2 . Thus we find the nine dimensional mapping

$$\begin{aligned}
 x_i &\mapsto x_{i+1}, & 0 \leq i < 5, \\
 x_4 &\mapsto x_5 = \frac{y_0 x_0}{y_0}, \\
 y_0 &\mapsto y_1, \\
 y_1 &\mapsto x_1 + y_0 - z_0 + z_2 - x_5, \\
 z_0 &\mapsto z_1, \\
 z_1 &\mapsto z_2 = \frac{x_5}{x_2} z_1.
 \end{aligned} \tag{D.1}$$

for which the staircase method does yield five functionally independent integrals. However, the mapping is not invertible and so the initial value problem is not well-posed. This example illustrates that in order to prove the well-posedness of an initial value problem one has to show the initial values can be updated in both directions.

REFERENCES

- [1] V.I. Arnold, *Dynamics of complexity of intersections*, Bol. Soc. Bras. Mat. **21** (1990), 1–10.
- [2] A.G. Akritas, E. K. Akritas and G. I. Malaschonok *Various Proofs of Sylvester's (Determinant) Identity* Mathematics and Computers in Simulation **42**, 585-593, 1996.
- [3] H.F. Baker, *Abelian Functions: Abel's Theorem and the Allied Theory of Theta Functions*, CUP, 1996.
- [4] E.H. Bareiss, *Sylvesters identity and multistep integer preserving Gaussian elimination*, Mathematics of Computation, **22**, 565-578, 1968.
- [5] H. Bateman and A. Erdelyi, *Orthogonal Polynomials*, in: *Handbook of Special Function*.
- [6] M.P. Bellon and C.-M. Viallet, *Algebraic entropy*, Comm. Math. Phys. **204** (1999), 425–437.
- [7] D. Bressoud, *Proofs and Confirmations: The Story of the Alternating Sign Matrix Conjecture*, CUP, 1999.
- [8] C. Brezinski, *Accélération de la Convergence en Analyse Numérique*, Lect. Notes in Math. **584** (Springer Verlag, 1977).
- [9] C. Brezinski, *Padé-Type Approximation and General Orthogonal Polynomials*, ISNM vol. **50**, Birkhäuser, Basel, 1980.
- [10] C. Brezinski, *A unified approach to various orthogonalities*, Annales de la Faculté des Sciences de Toulouse, Vol. **I**, n° 3 (1992), 277-292.
- [11] C. Brezinski, *Formal orthogonality on an algebraic curve*, Annals of Numerical Analysis **2** (1995) 21–33.
- [12] M. Bruschi, O. Ragnisco, P.M. Santini, and T.G. Zhang, *Integrable symplectic maps*, Physica D **49** (1991), 273–294.
- [13] L. Carlitz, *Some orthogonal polynomials related to elliptic functions*, Duke Math. J. **27** (1960) 443–460; *ibid.* II Arithmetic properties, Duke Math. J. **28** (1961), 107–124.
- [14] T.S. Chihara, *An introduction to orthogonal polynomials*, Gordon and Breach, New York, 1978.
- [15] A. Draux, *Polynômes Orthogonaux Formels, Applications*, LNM vol. 974, Springer-Verlag, Berlin, 1983.
- [16] G.Falqui and C.-M. Viallet, *Singularity, complexity and quasi-integrability of rational mappings*, Comm. Math. Phys. **154** (1993), 111–125.
- [17] W. B. Gragg, *The Padé table and it's relation to certain algorithm of numerical analysis*, (SIAM Review Vol. 14 1972) 1–62.
- [18] Grammaticos et al., *epsilon algorithms etc.*, Phys. Lett. **A**
- [19] F. Haggar, G.B. Byrnes, G.R.W. Quispel and H.W. Capel, *k-integrals and k-Lie symmetries in discrete dynamical systems*, Physica **233A** (1996) 379–394.
- [20] P. Henrici, *The quotient-difference algorithm*, in: *Further Contributions to the Solution of Simultaneous Linear Equations and the Determination of Eigenvalues*, Natl. Bur. of Standards Applied Mathematics Series **49**, US Government Printing Office (Washington, 1958). p. 23.
- [21] J. Hietarinta and C.-M. Viallet, *Singularity confinement and chaos in discrete systems*, Phys. Rev. Let. **81** (1998), 326–328.
- [22] J. Hietarinta and C.-M. Viallet, *Discrete Painlevé I and singularity confinement in projective space*, Solitons and Fractals **11** (2000), 29–32.
- [23] R. Hirota, *Nonlinear partial difference equations, I-III*, J.Phys.Soc.Japan **43** (1977) 1424–1433, 2074–2089.
- [24] D. Jackson, *Formal properties of orthogonal polynomials in two variables*, Duke Math. J. **2** (1936) 423–434.
- [25] Peter H. van der Kamp, *Initial value problems for lattice equations*, J. Phys. A: Math. Theor. **42** (2009) 404019.
- [26] Peter H. van der Kamp, *Growth of degrees of integrable mappings*, submitted to Journal of difference equations and applications (2009).
- [27] Peter H. van der Kamp and G.R.W. Quispel, *The staircase method*, In preparation.
- [28] G. Kowalewski, *Einführung in die Determinantentheorie*, Chelsea, New York, 1948.
- [29] H.L. Krall and I.M. Sheffer, *Orthogonal polynomials in two variables*, Ann. Mat. Pura Appl. (4) **76** (1967), 325–376.

- [30] I.M. Krichever and S.P. Novikov, *Holomorphic Fiberings and Nonlinear Equations*, Sov. Math. Dokl. 20 (1979) 650654; *Holomorphic Bundles over Algebraic Curves and Nonlinear Equations*, Russ. Math. Surv. 35 (1981) 5379.
- [31] J.S. Lomont and J. Brillhart, *Elliptic Polynomials*, (Chapman and Hall CRC, 2000).
- [32] G.I. Malaschonok, *Solution of a system of linear equations in an integral domain USSR J. Comp. Math. and Math. Phys.* **23**, 14971500, 1983 (in Russian).
- [33] G.I. Malaschonok, *System of linear equations over a commutative ring*, Academy of Sciences of Ukraine, Lvov, 1986 (in Russian).
- [34] B.M. McCoy and T.T. Wu *Nonlinear partial difference equations for the two-dimensional Ising model*, Phys. Rev. Lett. **45**, 675–678 (1980).
- [35] T. Muir, *The Theory of Determinants in the Historical Order*, Dover, New York, 1960.
- [36] M. Nieszporski and P.M. Santini, *The self-adjoint 5-point and 7-point difference operators, the associated Dirichlet problems, Darboux transformations and Lelievre formulae*, Glas. Math. J. (2005), **47**:133–147 CUP.
- [37] Y. Ohta, K. M. Tamizhmani, B. Grammaticos, and A. Ramani, *Singularity confinement and algebraic entropy: the case of the discrete Painlevé equations* Phys. Lett. A **262** (1999), 152–157.
- [38] V.G. Papageorgiou, F.W. Nijhoff and H.W. Capel. *Integrable mappings and nonlinear integrable lattice equations*, Phys. Lett. A **147** (1990) 106–114.
- [39] V. Papageorgiou, B. Grammaticos and A. Ramani, *Orthogonal polynomial approach to discrete Lax pairs for initial boundary-value problems of the QD algorithm*, Lett. Math. Phys. **34(2)** (1995), 91–101.
- [40] J.H.H. Perk, *Quadratic identities for Ising model correlations* Phys. Lett. A **79** (1980), 3-5.
- [41] G.R.W. Quispel, H.W. Capel, V.G. Papageorgiou and F.W. Nijhoff, *Integrable mappings derived from soliton equations*, Physica A **173**.
- [42] O. Rojas, Peter H. van der Kamp and G.R.W. Quispel, *Lax representations for integrable mappings*, In preparation.
- [43] P.E. Spicer and F.W. Nijhoff, *Polynomials in two variables, orthogonal over an elliptic curve*, In preparation.
- [44] P.K. Suetin, *Orthogonal polynomials in two variables*, (HarperCollins Publ., 1999).
- [45] M. Toda, *Theory of nonlinear lattices*, Springer-Verlag (Berlin, New York), 1981.
- [46] A.P. Veselov, *Integrable maps*, Russian Mathematical Surveys **46** (1991), 1–51.
- [47] P. Wynn, *The epsilon algorithm and operational formulas of numerical analysis*, Math. Comp. **15** (1961) 151–158.

KATHOLIEKE UNIVERSITEIT LEUVEN, DEPARTEMENT WISKUNDE, CELESTIJNENLAAN 200B B-3001 LEUVEN, BELGIUM

E-mail address: Paul.Spicer@wis.kuleuven.be

SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS LS2 9JT, U.K.

E-mail address: nijhoff@maths.leeds.ac.uk

DEPARTMENT OF MATHEMATICS AND STATISTICS, LA TROBE UNIVERSITY, VICTORIA 3086, AUSTRALIA

E-mail address: p.vanderkamp@latrobe.edu.au